



LABELS (PTY) LTD
Thru' Service We Sell

Date	<input type="checkbox"/>	24.08.18	Description	<input type="checkbox"/>	ZAR001/001-002
Customer	<input type="checkbox"/>	Fertex Group	Quantity	<input type="checkbox"/>	1 000
Order No.	<input type="checkbox"/>	Wendell	Size	<input type="checkbox"/>	135mm x 205mm Die
Contact	<input type="checkbox"/>	Wendell	Material	<input type="checkbox"/>	Dairyfilm HT Acrylic
Tel No.	<input type="checkbox"/>	N/A	Core Size	<input type="checkbox"/>	40
Fax No.	<input type="checkbox"/>	N/A	Labels p/roll	<input type="checkbox"/>	500
Email	<input type="checkbox"/>	fertexgroup@gmail.com	Roll Direction	<input type="checkbox"/>	Hand Applied
Barcode	<input type="checkbox"/>	N/A	Revision	<input type="checkbox"/>	6
Special Instructions	<input type="checkbox"/>	Design	Colours:	<input type="checkbox"/>	CMYK, Pant. 186C, Pant. 355C

PLEASE CHECK THE PROOF CAREFULLY.

Mark any corrections that are necessary and return. We are NOT responsible for errors not marked by you. WE CANNOT PROCESS YOUR ORDER WITHOUT A SIGNED PROOF. Proofs outstanding for two weeks, or orders cancelled, will be charged to your account.

Approved By: _____ **Date:** _____ **Alterations to be made ?** _____

205mm

DIRECTIONS FOR USE: Use only as recommended

SPRAY APPLICATION DIRECTIONS:

Crop Applications: (Agricultural Uses – For application to berry and small fruit crops, pome fruit, stone fruit, cereal grains, soybeans, and sunflowers.)

Use this product in areas to repel birds that congregate in large numbers, including these passerine birds: starlings, gulls, and rock dove and domestic pigeons, cliff swallow, house (English) sparrow, house finch, as well as these wild migratory waterfowl: geese, mute swan, and coot.

Ground Applications: This product must be applied using appropriate ground spray equipment. Withhold treatment until crop is susceptible to bird damage. Do not apply when crop is wet or rain is expected. Do not apply through any type of irrigation system. Product is best applied diluted while continuously agitated. For ground applications, 900 ml to 3 ℓ AVIAN CONTROL® per hectare with water at a ratio of 1 part product to 10-20 parts water or in sufficient water for the application equipment to be used. Apply at rates of 200 litres to 900 litres of spray mixture per hectare depending on application equipment and vegetation density. Apply the spray mixture so that the crop receives thorough coverage. This product may be applied up to the day of harvest (Pre-Harvest Interval = 0 days). Repeat treatment at 6-8 day intervals if significant bird damage begins to occur. Do not apply more than 35 litres per hectare per application.

Air Applications: (fixed wing and rotary aircraft): Apply the spray mixture so that the crop receives thorough coverage. Product is best applied diluted while continuously agitated. Apply 900 ml – 3 ℓ AVIAN CONTROL®. Dilute with water and apply at rates of 20 – 90 litres of spray mixture per hectare. This product may be applied up to the day of harvest (Pre-Harvest Interval = 0 days). Repeat treatment at 6-8 day intervals if significant bird damage begins to occur.

GROUP CODE 3 GROEP CODE

Active Ingredient/AktieweBestanddeel

Methyl Anthranilate (MA).....20%..... MetielAntranilate
Inert ingredients.....80%.....Ander bestanddele

NET Volume..... ℓ Nettovolume

Registered & Distributed By:



Fertex Group CC
(Reg. No. 2010/112400/23)
PO Box 135, Bergville, 3350, KZN, RSA
TEL: +27 (76) 900-8855
info@fertex.co.za, www.fertex.co.za

BATCH NO.:

MANUFACTURE DATE:

EXPIRY DATE:

IMPORTANT
Buffer water to PH 7
for optimal results.

Avian Control®

Bird Repellent

REG. NO. L10444 ACT No. 36 of 1947 / WET Nr. 36 van 1947



1ℓ 5ℓ 20ℓ 1000ℓ



CAUTION VERSIGTIG

135mm

205mm

Avian Control®

Bird Repellent

REG. NO. L10444 ACT No. 36 of 1947 / WET Nr. 36 van 1947

STORAGE & DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store only in original container, in a dry place inaccessible to children, pets and domestic animals. Store apart from pesticides, fertilizers, food or feed that may cause cross-contamination from odor. Keep container closed when not in use.

PESTICIDE DISPOSAL: Product or rinse water that cannot be used according to label instructions must be disposed of according to local and national regulations. Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING:

Non refillable Container (20 litres or less): Non-refillable container. Do not re-use or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Non refillable Container (greater than 20 litres): Non-refillable container. Do not reuse or refill this container. Offer for recycling, if available. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times.

Refillable Container: Refillable container. Refill this container with **AVIAN CONTROL®** only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the re-filler. To clean the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

WARRANTY

NOTICE: The directions for use of this product are believed to be adequate and must be followed carefully. To the extent consistent with applicable law, Avian Enterprises, LLC, Inc. does not make, nor does it authorize any agent or representative to make any oral or written warranty, guarantee, or representation, express or implied, concerning this material or the use there of beyond the statements of the label. To the extent consistent with applicable law, neither Avian Enterprises, LLC nor the seller shall be held responsible in any manner for any claim of personal injury or property damage or loss resulting from handling, storage, or use of this material not in accordance with the written directions. To the extent consistent with applicable law, the buyer assumes all risk and liability resulting from handling, storage, or use, or purchaser accepts and uses this material on these conditions. No other warranty exists.

CAUTION / VERSIGTIG

WARNINGS:

Handle with care. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes, and clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet. Store under lock and key, in a cool place. Store away from food and feeds. Toxic to fish, bees and moderately toxic to wildlife. Re-entry: Do not enter treated area for 4 hours or until spray deposit has dried unless wearing protective clothing. In case of poisoning a person, **CALL A DOCTOR** and in the case of an animal a **VETERINARIAN IMMEDIATELY AND MAKE THIS LABEL AVAILABLE TO HIM.**

Avian Control® is the registered trade name of / is die geregistreerde handelsnaam van Avian Enterprises Limited LLC, 2000 Pontiac Drive, Sylvan Lake, MI USA

PRECAUTIONS:

Do not inhale the spray mist. Wash contaminated clothing after use. Wash with soap and water after use and accidental contact. Do not eat, drink or smoke whilst mixing or applying the product or before washing hands and face after handling the product. Avoid drift of spray onto other crops, grazing, rivers, dams and areas not under treatment. Clean the applicator after use. Dispose of wash water where it will not contaminate crops, grazing, rivers, dams and boreholes. **TRIPLE RINSE** empty containers in the following manner: Invert the empty container over the spray or mixing tank and allow draining for at least 30 seconds after the flow has slowed down to a drip. Thereafter rinse the container three times with a volume of water equal to a minimum of 10 % of that of the container. Add the rinsing to the contents of the spray tank before destroying the container in the prescribed manner. Destroy the empty container by perforation and flattening and dispose of it in a safe way. Do not re-use the empty container for any other purpose. Prevent contamination of food, feeds, drinking water and eating utensils.

FIRST AID TREATMENT

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

Avian Control		Dosage per hectare
Label section	Subsection	
Crop Applications	Ground applications	900 ml to 3 ℓ per hectare
	Water rates per application	200 ℓ to 900 ℓ per hectare
	Maximum product application rate	35 ℓ per hectare
	Aerial applications	900 ml to 3 ℓ per hectare
	Water rates per application	20 ℓ to 90 ℓ per hectare

Crop Group	Pest	Commodities in Crop Group Approved for AVIAN CONTROL®
Pome Fruit	Cape white-eye (<i>Zosterops capensis</i>), Cape Sparrow (<i>Passer melanurus</i>), House sparrow (<i>Passer domesticus</i>) Red-winged Starling (<i>Onychognathus morio</i>), Common Starling (<i>Sturnus vulgaris</i>), White-backed mousebird (<i>Colius colius</i>), Red-faced Mousebird (<i>Urocolius indicus</i>), Speckled mousebird (<i>Colius striatus</i>) Cape Weaver (<i>Ploceus capensis</i>), Cape bulbul (<i>Pycnonotus capensis</i>), Blackkeyed bulbul (<i>Pycnonotus barbatus</i>), Red-eyed bulbul (<i>Pycnonotus nigricans</i>), Cape bulbul (<i>Pycnonotus capensis</i>)	Apple, pear, quince.
Stone Fruit		Apricots, cherries, sour cherries, sweet cherries, nectarines, peaches and plums.
Berry and Small Fruit		Blackberries, blueberries, boysenberries, cranberries, currants, dewberries, gooseberries, grapes, loganberries, raspberries, strawberries.
Cereal Grains	Red-billed quelea (<i>Quella quelea</i>), Southern Red Bishop (<i>Euplectes orix</i>), Cape weaver (<i>Ploceus capensis</i>), Southern masked weaver (<i>Ploceus velatus</i>), Cape sparrow (<i>Passer melanurus</i>) Streaky - headed Seedeater (<i>Crotophaga menelli</i>) Spur-winged Goose (<i>Plectropterus gambensis</i>), Egyptian goose (<i>Alopochen aegyptiaca</i> .)	Barley, maize, Oats, rice, rye, sorghum, sweetcorn, wheat.
Sunflowers		

135mm